Gravitation & Cosmology, Vol. 12 (2006), No. 2-3 (**46-47**), pp. 103–105
Proceedings of the International Conference on Gravitation, Cosmology, Astrophysics and Nonstationary Gas Dynamics,
Delicated to Prof. K.P.Staniukovich's 90th birthday, Moscow, 2-6 March 2006
© 2006 Russian Gravitational Society

arXiv:1105.6328v1 [gr-qc] 31 May 2011

EXACT STATIC SOLUTIONS FOR FLUID GRAVITATING BALLS IN HOMOGENEOUS COORDINATES

A.M.Baranov[1] **and R.V.Bikmurzin**[2]

Dep. of Theoretical Physics,Krasnoyarsk State University, 79 Svobodny Av., Krasnoyarsk, 660041, Russia
Received 22 April 2006

Two new classes of exact interior static solutions of the Einstein equations in homogeneous coordinates for a gravitating ball filled by a Pascal perfect fluid are obtained. Schwarzschild's interior solution of is a special case of these solutions.

1. Introduction

The choice of a coordinates' system can simplify the representation of the Einstein equations. A spherically symmetric metric interval in 4D space-time is usually written in of the curvature coordinates. In this paper, we make use of the homogeneous coordinates according to the Weyl remark [1] that the 3D spherical metric interval is conformally flat in such coordinates. So the static spherically symmetric metric interval for an interior gravitational field of a fluid stellar model with a general distribution of mass density is written as

$$ds^2 = A(r)dt^2 - B(r)\left(dr^2 + r^2(d\theta^2 + \sin^2\theta\, d\varphi^2)\right) =$$

$$= R_0^2(A(x)d\tau^2 - B(x)dl^2). \tag{1}$$

Here we introduce the dimensionless coordinates $\tau = t/R_0$; $x = r/R_0$; $0 \le x \le 1$, where t is the time variable; r is the radial variable; $\theta; \varphi$ are angular variables; R_0 is a stellar exterior radius; $dl^2 = (dx^2 + x^2(d\theta^2 + \sin^2\theta\, d\varphi^2))$; the velocity of light $c = 1$; $A(x)$ and $B(x)$ are metric functions.

2. The first class of solutions

The Einstein equations for the energy-momentum tensor of Pascal's perfect fluid

$$T_{\alpha\beta} = (\mu(x) + p(x))\, U_\alpha U_\beta - p(x)\, g_{\alpha\beta} \tag{2}$$

are transformed to three equations for $A(x)$ and $B(x)$ functions. The 4-velocity in a comoving frame of reference is equal to $U_\alpha = \delta_\alpha^0 \sqrt{A(x)R_0^2}$.

In the case with $\mu = const$ (a homogeneous distribution of mass), one of these equations may be rewritten as Emden's equation [2]

$$\frac{d^2 b(x)}{dx^2} + \frac{2}{x}\frac{db(x)}{dx} + b(x)^5 = 0. \tag{3}$$

From this equation we can find the function $B(x)$ and then, from other gravitational equations, we obtain function $A(x)$ [3]

$$B_{Sch.}(x) = \frac{C_1}{(C_2 + x^2)^2},$$

$$A_{Sch.}(x) = \left[C_4 - \frac{C_3}{(C_2 + x^2)}\right]^2. \tag{4}$$

Thus we have obtained Schwarzschild's interior solution in homogeneous coordinates. The Weyl tensor is equal to zero for such exact solution which describes the conformally flat 4D space-time. The junction conditions on the stellar surface give connections of constants with compactness $\eta = 2m/R_0$, where m is Schwarzschild's mass and $0 \le \eta < 1$.

When $\mu(x)$ is a function of x, we take the function $B(x)$ as a generalization of the case with a homogeneous distribution of mass density in the form

$$B_{in}(x) = \frac{C_1}{(C_2 + x^2)^k} = C_1 Y(x)^{-k}, \tag{5}$$

where $Y(x) = \left(C_2 + x^2\right)$ and $0 \le k \le 2$, then we can get functions $A_{in}(x)$, $\mu(x)$ and $p(x)$:

$$A_{in}(x) = \left[\frac{C_4 Y(x)^\Lambda - C_3}{Y(x)^{(k+\Lambda-1)/2}}\right]^2; \tag{6}$$

$$\mu(x) = \frac{kY(x)^{k-2}}{8\pi R_0^2 C_1}\left[(2-k)Y(x) + (k+4)C_2\right]; \tag{7}$$

$$p(x) = \frac{Y(x)^{k-2}}{4\pi R_0^2 C_1}\left[Y(x) + 7kC_2 + \frac{C_3 + C_4 Y(x)^\Lambda}{C_3 - C_4 Y(x)^\Lambda}\times\right.$$

$$\left.\times\Lambda\big((k-1)Y(x) - kC_2\big)\right] - 3\mu(x), \tag{8}$$

where $\Lambda = \sqrt{2k^2 - 4k + 1} \ge 0$ and the parameter k can have values $k = \left(0; 1 - 1/\sqrt{2}\right] \cup \left[1 + 1/\sqrt{2}; 2\right]$.

In this case, the functions $B(x)$ and $\mu(x)$ are chosen as nonsingular functions for the stellar source. We also take the pressure equal to zero on the stellar surface.

---

[1] e-mail: bam@lan.kras.ru

[2] e-mail: Rustam_B_V@mail.ru

So the metric interval (1) may be rewritten as

$$ds^2 = \frac{R_0^2 C_1}{Y(x)^k}\left[\frac{Y(x)^{1-\Lambda}}{C_1}(C_4Y(x)^{\Lambda}-C_3)^2 d\tau^2 - dl^2\right] \tag{9}$$

Now we must find the constants C_1, C_2, C_3 and C_4 for a final result.

As an exterior solution, we take the Schwarzschild exterior solution in homogeneous dimensionless coordinates (see, e.g., [4])

$$A_{ext}(x) = \left(\frac{4x-\eta}{4x+\eta}\right)^2, \quad B_{ext}(x) = \left(1+\frac{\eta}{4x}\right)^4. \tag{10}$$

Using the smooth junction conditions on the 2D stellar surface with the exterior solution in the form (10) in homogeneous coordinates, we have

$$C_1 = \left(1+\frac{\eta}{4}\right)^4\left[\frac{k(4+\eta)}{2\eta}\right]^k; \tag{11}$$

$$C_2 = \frac{k(4+\eta)}{2\eta} - 1; \tag{12}$$

$$C_3 = \frac{k}{4\eta\Lambda}\Big(4k-(\eta-4)(k-1-\Lambda)\Big)\left[\frac{k(4+\eta)}{2\eta}\right]^{(k-3+\Lambda)/2}; \tag{13}$$

$$C_4 = \frac{k}{4\eta\Lambda}\Big(4k-(\eta-4)(k-1+\Lambda)\Big)\left[\frac{k(4+\eta)}{2\eta}\right]^{(k-3-\Lambda)/2}. \tag{14}$$

From a requirement $C_2 > 0$ it follows

$$\frac{k(\eta+4)}{2\eta} - 1 > 0 \;\Rightarrow\; k > \frac{2\eta}{\eta+4} \;\text{ or }\; \eta < \frac{4k}{2-k}. \tag{15}$$

If the parameter k is in the range $\left(0; 1-1/\sqrt{2}\right]$, then the compactness η has values in the interval $\left[0;\, 4(\sqrt{2}-1)/(\sqrt{2}+1)\right) \approx [0;\, 0.686)$. If $k > 1+1/\sqrt{2}$, then parameter η is arbitrary.

As a result, all functions of the first-class solutions may be rewritten in a clear form:

$$B_{in}(x) = \left(1+\frac{\eta}{4}\right)^4\left[1-\frac{2\eta}{k}\frac{(1-x^2)}{\eta+4}\right]^{-k} = \left(1+\frac{\eta}{4}\right)^4 \Psi(x)^{-k}; \tag{16}$$

$$A_{in}(x) = \left[\frac{(\eta-4)(k-1-\Lambda)-4k}{2\Lambda(\eta+4)}\Psi(x)^{(1-k\Lambda)/2} \times \left(1-\xi\Psi(x)^{\Lambda}\right)\right]^2; \tag{17}$$

$$\mu(x) = \frac{128\eta}{\pi R_0^2 k}\frac{\Psi(x)^{k-2}}{(\eta+4)^6}\left[12k+\eta(k-2)(3-x^2)\right], \tag{18}$$

where the following notations are used:

$$\xi = \frac{(\eta-4)(k-1+\Lambda)-4k}{(\eta-4)(k-1-\Lambda)-4k};$$

$$\Psi(x) = \left[1-\frac{2\eta}{k}\frac{(1-x^2)}{(\eta+4)}\right]. \tag{19}$$



Figure 1. Behavior of the metric functions $A(x) = g_{00}(x)$ and $B(x) = g_{11}(x)$ of the complete solutions belonging to the first and second classes with $\eta = 0.2$ and $k = 0.1003$. The number 0 corresponds to Schwarzschild's interior solution. The numbers 1 and 2 correspond to the first and second classes of solutions.

3. The second class of solutions

Now we shall generalize the function $A(x)$ from (4) by a similar way. The function $A(x)$ is defined as

$$A_{in}(x) = C_1(C_2 - x^2) = C_1 Y(x)^k. \tag{20}$$

Then the Einstein equations give

$$B_{in}(x) = \left[\frac{Y(x)^{(\Lambda-1-k)/2}}{C_4Y(x)^{\Lambda}-C_3}\right]^2; \tag{21}$$

$$p\,(x) = \frac{Y(x)^{k-\Lambda-1}}{8\pi R_0^2}\Big[\Big(k^2\big(Y(x)-C_2\big)-2C_2(1+\Lambda)\Big)\times \times\big(C_3+C_4Y(x)^{\Lambda}\big)^2 + \Big(Y(x)(1+k^2)+C_2(1+\Lambda-k^2)\Big)\times \times 4C_3C_4Y(x)^{\Lambda} + 4C_2C_3^2\Lambda\Big]; \tag{22}$$

$$\mu(x) = \frac{k\big(C_3-C_4Y(x)^{\Lambda}\big)}{4\pi R_0^2 Y(x)^{1+\Lambda-k}}\Big[\Big(3C_2-\Lambda\big(Y(x)-C_2\big)\Big)\times \times\big(C_3-C_4Y(x)^{\Lambda}\big)+2\Lambda C_3(Y(x)-C_2)\Big] - 3p(x), \tag{23}$$

where $\Lambda = \sqrt{1+2k^2}$.

Under the smooth junction conditions with Schwarzschil's exterior solution, the new constants are

$$C_1 = \left(\frac{4-\eta}{4+\eta}\right)^2\left[\frac{8\eta}{k(16-\eta^2)}\right]^k; \tag{24}$$



Figure 2. Behavior of the mass density $\mu(x)$ of the total solutions belonging to the first and the second classes with $\eta = 0.2$, $k = 0.1003$ and a central mass density $\mu_0 = const.$ The number 0 corresponds to Schwarzschils's interior solution. The numbers 1 and 2 correspond to the first and the second classes of solutions.

$$C_2 = \frac{k(16-\eta^2)}{8\eta} - 1; \tag{25}$$

$$C_3 = \frac{2\big(4(\Lambda+1)+k\eta\big)}{(\eta+4)^2\Lambda}\left[\frac{8\eta}{k(16-\eta^2)}\right]^{(1+k-\Lambda)/2}; \tag{26}$$

$$C_4 = \frac{2\big(4(\Lambda-1)-k\eta\big)}{(\eta+4)^2\Lambda}\left[\frac{8\eta}{k(16-\eta^2)}\right]^{(1+k+\Lambda)/2}. \tag{27}$$

Finally, we have bounds for the parameter k

$$\frac{80+40\eta-7\eta^2+\chi}{\eta^3-10\eta^2-64\eta+480} \geq \frac{k}{2} \geq \frac{4\eta}{(16-\eta^2)}, \tag{28}$$

where χ is

$$\chi = \sqrt{17\eta^4 - 240\eta^3 + 2528\eta^2 - 8960\eta + 6400}. \tag{29}$$

4. Summary

The two classes of interior exact solutions of Einstein's equations, found in the paper, describe the gravitational field of a ball filled with a Pascal perfect fluid. These solutions correspond to a static stellar model which is a generalization of Schwarzschild's interior solution. Limiting conditions for the parameters have been found. The results of this study have been applied to a neutron star model with $\eta = 0.2$ and the parameter $k = 0.1003$. Plots of the metric functions and the mass density have been constructed.

References

[1] L.P.Eisenhart, "Riemannian Geometry", Princeton Univ. Press, Princeton, 1949.

[2] G.Sansone, "Equazioni Differenziali Nel Campo Reale", Bologna, Partre Seconda, 1949.

[3] A.M.Baranov, R.V.Bikmurzin, *Abstr. of Int. Seminar "Volga-16"*, Kazan Univ., Kazan,2004, p.24-25.

[4] N.M.Mitskievich, "Physical Fields in General Relativity", Nauka, Moscow, 1969 (in Russian).